The geologic record of hydration and dehydration in the subducting slab: Epidote minerals record alteration and metamorphism before and during subduction


Peter C. Lindquist¹, Cailey B. Condit¹, William F. Hoover¹, Victor E. Guevara²

¹University of Washington, ²Amherst College


Contact: plindqui@uw.edu



The fingerprint of metamorphic dehydration in subduction zones





The breakdown of hydrous minerals like lawsonite and chlorite are expected to release significant volumes of water from subducting basalts. These fluids influence the rheology of plate interface (e.g., promoting slow slip and tremor). Epidote is a common product of the predicted dehydration reactions.



We analyze epidote in metabasalts from the Catalina Schist to (1) explore the use of epidote as a tracer of prograde subduction processes and (2) look for evidence of dehdration reactions in the subducted rock record.

The Catalina Schist represents an exhumed Cretaceous subduction complex, composed of tectonic slices of oceanic crust that were subducted and underplated to North America. The epidote-amphibolite facies unit experienced peak P-T conditions of 0.8–1.3 GPa and 500–560°C, representative of conditions at ~35 km depth in modern warm subduction zones, around 105 Ma.






Distinct from interpillow epidosite networks, we interpret wispy, epirote-rich vein-like features as forming during prograde metamorphism during subduction.


Epidote-rich networks preserve early metamorphism

"Ghost grains" in the cores of epidote crystals likely represent replacement of pumpellyite based on crystal shapes and geochemistry. REE patterns in these ghost grains are comparable to those measured in pumpellyite from other metabasalts.



1(2)


While metamorphic epidote-rich zones lack discrete vein-like boundaries, elemental X-ray maps reveal discrete Ti-poor zones within these. These may represent zones of transient porosity produced during pumpellyite breakdown.



Metamorphic epidote-rich zones may represent self-organized dehydration features that developed as a result of intial compositional heterogeneity.

Porphyroblastic epidote records pro- and retrograde metamorphism

While metamorphic epidote networks preserve ~intial growth, epidote porphyroblasts react with surrounding matrix and exhibit complex textures and compositions that track progressive metamorphism




identify two generations of amphibole and epidote:


> Both hornblende and actinolite are present in the matrix and within porphyroblast structures. Hornblende is commonly rimmed by actinolite.

cores of more Fe-rich epidote (Ep₂) are embedded within more Al-rich epidote (Ep_3).

Hornblende is embayed by the



Pairing these textures and mineral compositions with phase equilibrium modeling with PerpleX constrains a range of P-T conditions for the growth of the two generations of minerals.




The abundance of epidote is predicted to decrease with increase temperature, which helps explain the presence of Ep₂ as relic cores that are then overgrown by Ep₃.

Epidote 2 therefore represents the composition of epidote at ~peak P-T conditions, while growth of more Al-rich epidote 3 with actinolite suggests retrograde cooling.

• REE patterns for epidote porphyroblasts (both Ep₂ and Ep₃) are enriched in LREEs relative to epidote-rich networks (Ep₄).

> • With progressive metamorphism, epidote porphyroblasts likely preferentially incorporated more LREE from other reacting phases while the epidote-rich networks do not communicate as much










Records of seafloor alteration

Pillow and hyaloclastite breccia textures are preserved in parts of the epidote amphibolite metabasalts.


Networks of interpillow material are commonly epidote rich and we interpret these as having been altered to epidosite.

Strongly negative Ce anomalies in from these zones are interaction with seawater.




Jurassic seawater ⁸⁷Sr/⁸⁶Sr Bulk-rock Sr isotope measurements provide another indication of intial seawater-rock range of initial ⁸⁷Sr/⁸⁶Sr for ep. interactions, and amphibolites suggest potentially large water/rock ratios prior to subduction. ~MORB 87 Sr/865

epidosites record hydration of oceanic crust prior to subduction




Pumpellyite to epidote dehydration reactions result in a pulse of water release around 250-300°C



H₂O fluxes

North America



epidote growth

reaches a peak

around 400°C

Phase equilibrium modeling also predicts the conditions at underplating at ep. nphibolite conditions

This transition is associated with a release of ~94 kg H₂O/m³ of basalt, which, to first order, could result in fluxes of 10^4 – 10^5 kg H₂O/m²/Myr in the region of such dehydration in a generic refrigeration following

underplating restabilizes more epidote (Ep₃) underplating at 95 Ma

North America

with increasing T to peak

conditions the volume %

Retrograde Ep2 sonite blueschist unit

Si-content of hornblende. The

Ep2 represents near-peak

epidote compositions

of actinolite and albite.

with the matrix.





