


Inter-event strain localization modulated by background // stresses across the Natron Basin, East African Rift









ETH zürich Iván Navarrete¹, Jean-Arthur Olive¹, Eric Calais¹, Manon Dalaison², Albert de Montserrat³

Introduction

- Plate separation at continental rifts is accommodated by tectonic and magmatic processes.
- On short time scales, rifting manifests as discrete events of fault slip and magmatic intrusion.
- Such events relieve extensional stresses built-up at the axis over 10's to 100's years.
- Here, we focus on the inter-event localization of deformation in the Natron Basin

Key Question:


Does long-term (finite) deformation influence short-term stress build-up?


Study Area: Natron Basin - Part of the eastern branch of the East African Rift - Experienced a major tectono-magmatic event in 2007. (Calais et al 2008) GNSS horizontal velocity profile across the rift reveals a gradual transition zone between the Tanzanian craton and the Somalian plate, with ~2 mm/year of extension distributed over a ~ 100 km stretching zone.

So, what controls the width of this

inter-event stretching zone?

Stretching Zone: Prior Models Dislocation (back-slip) Model Natron border fault Requires localized shear zone deep within the lower crust. Inconsistent with weak, magma-rich lower crust? Elastic Necking model Produces no stretching at the axis. Strain localization would require weaker elastic moduli at the axis





Next Steps:

- Usage of pseudo-transient numerical methods, where fundamental conservation equations are formulated in terms of strain increments rather than strain rate.
- Analyze the impact of the brittle layer thickness in the width of the stretching zone.

Take Home Messages

- The width of the stretching zone in the Natron Basin can be explained by the accumulation of heterogeneous stresses due to past deformation, particularly fault block bending, rather than by continuous slip on deep faults.
- A 2D thermo-mechanical model shows that even in the absence of deep fault slip, localized deformation occurs due to preexisting stress fields, producing a sigmoidal velocity profile.
- This modeling challenges classic elastic back-slip interpretations and suggests that **long-term stress** accumulation significantly controls short-term deformation in magma-influenced continental rifts like the

Acknowledgement

This work was funded by the Marie Skłodowska-Curie Actions under the European Union's Horizon research and innovation programme through the TALENTS Doctoral **Network** . I also gratefully acknowledge the **GPU4GEO** group for their invaluable support and the development of tools that greatly facilitated this research.