

Observation of the Changing Earth from Space

Hartmut Grassl

Max Planck Institute for Meteorology, Hamburg

European Geosciences Union – General Assembly

Lecture at the
**Geoscience Information for Teachers (GIFT)
Workshop**


Austria Centre Vienna, 20 April 2009



Max-Planck-Institut
für Meteorologie

Carbon cycle

Source: IPCC AR4-WGI, 2007



Earth Observation from Space Contributes to

- Detection and Exploitation of Windows of Predictability
- Warning of Geohazards and Mitigation of their Impacts
- Control of Compliance with International Law

Earth Observation from Space is the Backbone of Services

- established for weather forecasting
- in build-up phase for
 - oceanography
 - disaster prevention and mitigation
 - chemical weather forecast



Climate Data from Satellites (GCOS)

Source: GCOS IP 2004

Table 1. Essential Climate Variables that are both currently feasible for global implementation and have a high impact on UNFCCC requirements.


Domain	Essential Climate Variables
Atmospheric (over land, sea and ice)	<p>Surface: Air temperature, Precipitation, Air pressure, Surface radiation budget, Wind speed and direction, Water vapour.</p> <p>Upper-air: Earth radiation budget (including solar irradiance), Upper-air temperature (including MSU radiances), Wind speed and direction, Water vapour, Cloud properties.</p> <p>Composition: Carbon dioxide, Methane, Ozone, Other long-lived greenhouse gases, Aerosol properties.</p>
Oceanic	<p>Surface: Sea-surface temperature, Sea-surface salinity, Sea level, Sea state, Sea ice, Current, Ocean colour (for biological activity), Carbon dioxide partial pressure.</p> <p>Sub-surface: Temperature, Salinity, Current, Nutrients, Carbon, Ocean tracers, Phytoplankton.</p>
Terrestrial	<p>River discharge, Water use, Ground water, Lake levels, Snow cover, Glaciers and ice caps, Permafrost and seasonally-frozen ground, Albedo, Land cover (including vegetation type), Fraction of absorbed photosynthetically active radiation (fAPAR), Leaf area index (LAI), Biomass, Fire disturbance.</p>

- variable depends on satellite observations



Impact of Satellite Data on Forecast Skill


Source: The Changing Earth (SP-1304, ESA, 2006)



Increase in anomaly correlation of 500hPa height forecasts during recent decades is to a large extent due to the assimilation of satellite data

HOAPS-G Freshwater Flux

HOAPS-G Freshwater-Flux JAN1988-DEC2005




Trend Analyses with Satellites

Variable	Start of Analysis	Observed Trend
Sea Level	1992	+ 3.2 ± 0.5 mm/a
SST	1991	+ 0.13 ± 0.03 °C/10a
ozone column	1978	intensification of ozone hole until 2006
nitrogen dioxide	1995	+ 50% over China
cloud albedo	1981	-2% over Europe due to pollution abatement policies and collapse of Soviet Union
solar output	1979	no trend



Observed Global Sea Level Rise

Source: The Changing Earth (SP-1304, ESA, 2006)





sea level rise derived from several satellite altimeters



Cloud Reflectance over Europe around 1990


Source: H. Grassl / O. Krüger, 2002



Decrease of cloud reflectance (unit: %) in thousandth from 1985-89 to 1996-99 for different mean reflectance classes as derived from AVHRR channel 2 satellite measurements over parts of Central Europe for winter(left) and summer (right) months. This is interpreted as a net reaction to less aerosol and soot emission after the breakdown of the eastern European industry production around 1990.

Cloud Reflectance and Ship Emissions




Source: A. Devasthale, 2006



Trend observed in monthly averaged cloud albedo (upper panel) and CTTs (lower panel) between May 97 and Aug. 02 for 3 harbour areas (R1, R3), the English Channel (R2) and a relatively remote ocean region (R4) clearly confirms the influence of increasing ship emissions.

In addition to an albedo increase the reduction of cloud top temperature hints at more intense convective activity in aerosol laden clouds.

Understanding the Earth System




Increase in NO₂-concentration by 50%

Courtesy of John Burrows, Univ. Bremen, D

Stability of the Solar Constant

Source: PMOD/WRC in Davos



Challenges to be answered soon

- Source and sink distribution of CO, CH₄ and CO₂ by inverse modelling using data from ENVISAT, METOP
- Trends of cloudiness by better intercalibration of satellite series
- Phytoplankton time series through combination of CZCS, SeaWiFS, MODIS and MERIS



Did Earth Observation Influence International Law?

- yes, for the Montreal Protocol
- yes in parts, for EU climate policy if EU council decisions of March 2007 are implemented
- hopefully, for the follow-up to the Kyoto Protocol



Do we have a balanced European system of Earth Observation?

Component	Institution	Purpose
Earth Explorer Core and Opportunity Missions	ESA-EOEP	New observations with new technologies for understanding the Earth System
Operational Meteorological Satellites (built by ESA)	EUMETSAT	Improved weather forecasts Fundamental data for GMES services
Sentinel de la Terre (GMES operational satellite series)	European Commission + ESA	New operational services within GMES as part of GEOSS

Answer: Yes, soon



Thank you!

